

Covariant treatment of flavour oscillations

Subhendra Mohanty

Physical Research Laboratory, Navrangpura, Ahmedabad 380009, INDIA

February 1, 2008

Abstract

We perform a covariant wavepacket analysis of neutrino oscillations taking into account the lifetime of the neutrino production process. We find that flavour oscillations in space are washed out when the neutrinos are produced from long lived resonances - and what may be observed in appearance/disappearance experiments is a uniform conversion probability independent of distance. The lifetime of the resonance which produces the neutrinos acts as the effective baseline of the experiment. For this reason the LSND experiment where neutrinos are produced from muon decay has two orders of magnitude more sensitivity to neutrino mass square difference than other experiments where the neutrinos are produced from pion or kaon decays. We also show that there are no EPR type oscillations of the secondary decay particles.

1 Introduction

There are two different formulas which describe flavour oscillations in spacetime; the one applied to neutrino oscillations [1] is derived in the high energy regime and the other applied to kaon oscillations [2] is valid in the non-relativistic limit. From the derivations of these formulas it is not clear what formula is applicable at intermediate energies (for example for kaon oscillations from stopped protons where $(m^2/E^2) \sim 25\%$). A covariant derivation of a flavour oscillation formula which would be valid at both high and low energies would be of interest from the conceptual as well as the experimental point of view.

Kayser and Stodolsky [3] have advocated a covariant generalisation of the non-relativistic phase factor $\exp(-imt)$ by replacing the absolute time t by the Lorenz invariant proper time $s = (t^2 - x^2)^{1/2}$. In the lab frame the covariant expression for the phase factor is $\exp - im_i s_i = \exp(-im_i^2 t/E_i)$. The phase difference between two mass eigenstates is $\Delta(m^2/E) \simeq (\Delta m^2/E)$ which is twice the phase difference of the standard formula [1]. This result has prompted the claim [4] that kaon oscillations in Φ factory will have oscillation length which is half of what is given by the standard formula. This claim has been refuted by [5] who take the view that the interference phase difference should not be evaluated at different space-time points but should be evaluated at

the average spacetime interval and taking the phase difference to be $(m_1 - m_2)(s_1 + s_2)/2$ instead of $(m_1 s_1 - m_2 s_2)$ they recover the standard formula. Other methods of showing that an extra factor of two does not appear in the kaon oscillations formula have been discussed in [6]. A covariant derivation of the neutrino oscillation formula has been given by Grimus and Stockinger [7] who treat the entire process of neutrino production, propagation and detection as a single Feynman diagram. They show that on taking the large distance limit of the neutrino propagator the scattering cross section shows a space-time oscillatory behaviour and the oscillation length is identical to that given by the standard formula [1]. G-S assume the initial states to be plane waves therefore the concept of a coherence length [8] does not emerge in their formulation.

The oscillation phenomenon occurs because particles are produced and detected as weak interaction eigenstates (the neutrino states ν_e, ν_μ and ν_τ or the Kaon states $K^0, \bar{K}^0$ etc) but the propagators are diagonal in the mass eigenstates (the neutrino mass eigenstates $\nu_i, i = 1 - 3$ or the Kaon mass eigenstates K_L, K_S). The probability amplitude for oscillations of a gauge eigenstate ($|\alpha\rangle$) to another ($|\beta\rangle$) is a lot of a gauge eigenstate ($|\alpha\rangle$) to another ($|\beta\rangle$) is a linear superposition of the propagation amplitude of the mass eigenstates ($|i\rangle$).

$$\mathcal{A}(\alpha \rightarrow \beta; t) = \sum_i \langle \beta | i \rangle \langle i | e^{-iHt} | \alpha \rangle \quad (1.1)$$

In relativistic field theory the propagation amplitude of the mass eigenstates $\langle i | e^{-iHt} | i \rangle$ can be identified with the Feynman propagator. At large time-like spacetime separation, the Feynman propagator (for both fermions and bosons) has the form,

$$K(m_i, t) \simeq \left(\frac{m_i}{2\pi i s_i} \right)^{3/2} e^{-im_i s_i} \quad (1.2)$$

where s_i is the invariant space-time interval propagated by the ν_i mass eigenstate. In the lab frame the phase factor of the amplitude (1.2) is given by $\exp\left(\frac{-im_i^2 t}{E_i}\right)$. In this paper we derive the oscillation amplitude by evaluating the Feynman propagator of the neutrinos in the large time-like asymptotic limit. The asymptotic propagator of a position eigenstate has the form $K(x, t; m_i) \simeq (m_i/2\pi i s_i)^{3/2} \exp(-im_i s_i)$ where s_i is the spacetime interval propagated by the m_i mass eigenstate. We derive the same expression in the path integral method. Using the path integral method we show that for large distance propagation the forward in time paths are equivalent to on-shell particles.

Another aspect of flavour oscillations on which there is no consensus of opinion is the question of the energy momentum of a flavour eigenstate. A flavour eigenstate propagates as a linear combination of different mass eigenstates which are on-shell. Some authors have argued that the different mass eigenstates have common energy but different momenta $P_i = \sqrt{E^2 - m_i^2}$ while others have assumed that the states have same momenta but different energy $E_i = \sqrt{P^2 + m_i^2}$. We show here that different mass eigenstates in a linear combination of states have neither energy nor momentum in common. For long distance propagation, the energy and momenta of a given mass eigenstate are not independent however but are related by the mass shell condition. We use these results to show that there are no EPR type associated particle oscillations as has been claimed in literature [4]. This result is in agreement with the conclusions of Lowe et al. [5] and Dolgov et al. [6]. Our proof however is more direct in that we show that although the energy and momenta of the muons depend on the outgoing neutrino mass, the phase difference of the muon states corresponding to different neutrino masses cancels exactly and there is no spacetime dependence of the muon probability distribution.

If the initial wavefunctions of the propagating particles were strictly delta functions in spacetime then no interference between different mass eigenstates can take place. We therefore generalise the delta function propagators to propagators of gaussian wavepackets. The interference term as a function of distance is obtained by taking the time-overlap of different mass eigenstate propagators. The expression for flavour conversion probability (for say two flavours with mixing angle θ) as a function of distance X turns out to be,

$$P(\alpha \rightarrow \beta; X) = \frac{1}{2} \sin^2 2\theta (1 - \cos(\frac{\Delta m^2}{2P} X)) e^{-A} \quad (1.3)$$

Therefore extra factor of two which came from naively subtracting the phases of plane wave propagators [4] goes away in the wavepacket averaging and the standard expression for the oscillations length is recovered. This is in agreement with [5]-[6]. The covariant wavepacket treatment introduces a new contribution to the exponential factor A when particles are produced from long lived resonances (for example for neutrinos from muon decay as opposed to neutrinos from Z decay). At distances smaller than the coherence length , the exponential suppression factor is,

$$A = (\frac{\Delta m^2}{2\sqrt{2}E} \tau)^2 \quad (1.4)$$

where τ is the lifetime of the resonance which produces the particle which undergo flavour oscillations. If the uncertainty in position of the initial particle $v\tau$ is larger than the detector distance X , the exponential term washes out the oscillations in (1.3). In neutrino experiments where the source is pions, kaons, muons or nuclear fission, the spatial oscillations of the conversion probability cannot be observed. What can be observed in these experiments is a constant (distance independent) conversion probability. The conversion probability is sensitive to values of $\Delta m^2 \simeq (2\sqrt{2}E/\tau)$, which means that the experimental bound on Δm^2 is lower with longer -lifetime sources. For this reason the LSND experiment [9] which uses neutrinos from muon decay ($\tau_\mu = 2.19 \times 10^{-6}s$) is sensitive to two orders of magnitude lower neutrino mass square difference compared to other accelerator experiments like BNL-E776 [10], Karmen [11] and CCFR [12] which use neutrinos from pion and kaon decays ($\tau \sim 10^{-8}sec$). We fit the experimental data from LSND along with BNL-E776, Karmen, CCFR and Bugey [13] experiments with the conversion probability formula (1.3) and plot the allowed range for the mass difference and mixing angle. We find that there is a large region of LSND allowed band which is not ruled out by the other experiments. The region of Δm^2 and $\sin^2 2\theta$ which is allowed by LSND and all other experiments is $2.2 \times 10^{-3} eV^2 < \Delta m^2 < 6.3 eV^2$ and $0.002 < \sin^2 2\theta < 0.15$. The lower bound on allowed Δm^2 is two orders lower than what is obtained by fitting the LSND data with the standard oscillation formula (equation (1.3) with A set to zero).

2 Field theoretic derivation of phase factor

In relativistic field theory the propagation amplitude of the mass eigenstates $\langle i|e^{-iHt}|i \rangle$ can be identified with the Feynman propagator $S_F(x_f - x_i, m_i) = \langle T \nu_i(x_i) \bar{\nu}_i(x_f) \rangle$. The Feynman propagator in position space is ,

$$K(x, m_i) = -i \int \frac{d^4 p}{(2\pi)^4} \frac{(i \not{p} + m_i)}{p^2 - m_i^2 - i\epsilon} e^{-ip \cdot x}. \quad (2.1)$$

This integration can be done by expressing the denominator as an exponential,

$$\frac{-i}{p^2 - m_i^2 - i\epsilon} = \int_0^\infty d\alpha \exp \{i\alpha(p^2 - m_i^2 - i\epsilon)\} \quad (2.2)$$

and integrating over the resulting Gaussian in p . The remaining integral over α ,

$$\int_0^\infty d\alpha \alpha^{-2} \exp \left\{ i\alpha m_i^2 + i \frac{(t^2 - |\mathbf{x}|^2)}{4\alpha} \right\} \quad (2.3)$$

can be performed by substituting $s = (t^2 - |\mathbf{x}|^2)^{1/2}$, $z = ims$ and $\eta = 2(\alpha m/s)$ and making use of the integral formula [14] for the Bessel function

$$\frac{1}{2} \int_0^\infty d\eta \eta^{-(\nu+1)} \exp -\frac{z}{2} \left(\eta + \frac{1}{\eta} \right) = K_\nu(z). \quad (2.4)$$

The resulting expression for the Feynman propagator (2.1) is

$$K(x, m_i) = \left(\frac{i}{4\pi^2} \right) (i \not{\partial} + m_i) \left(\frac{m_i}{s_i} \right) K_1(im_i s_i) \quad (2.5)$$

where $s_i = (t^2 - \mathbf{x}_i^2)^{1/2}$ is the invariant spacetime interval propagated by the ν_i mass eigenstate. If this interval is large ($s \gg m_i^{-1}$) and time-like ($t \geq |\mathbf{x}|$) then we can use the asymptotic expansion of the Bessel function

$$K_1(ims) \simeq \left(\frac{2}{\pi ims} \right)^{1/2} e^{-ims} \quad (2.6)$$

to obtain from (2.5) the expression for the propagation amplitude at large time-like separation

$$K(x, t; m_i) = \left(\frac{m}{2\pi i \sqrt{(t_f - t)^2 - (\mathbf{x}_f - \mathbf{x}_i)^2}} \right)^{3/2} \exp \{ -im \sqrt{(t_f - t)^2 - (\mathbf{x}_f - \mathbf{x}_i)^2} \} \quad (2.7)$$

The phase factor of the amplitude (2.7) is Lorentz-invariant and can be written in terms of the neutrino energy E and the time of flight t as measured from the lab frame as $-im_i s_i = -im_i (1 - v_i^2)^{1/2} t = -i(m_i^2 / E_i) t$.

An extra factor of two appears on subtracting the phases at different spacetime points. A flavour eigenstates neutrino or kaon is observed at a single spacetime point and one should therefore compute the phase difference at the overlap of the two mass eigenstate wavepackets. This averaging over spacetime is done formally by considering the propagators of gaussian wavefunctions as opposed to plane waves, and the standard expression for the oscillation length is recovered.

3 Conservation laws for long distance propagators

A flavour eigenstate propagates as a linear combination of different mass eigenstates which are on-shell. The phase difference is obtained by some authors by assuming that the different mass eigenstates have common energy, and by some by assuming that they have a common momentum [16]. In this section we show that the linear combination of different mass eigenstates have neither the same energy nor the same momentum. We show that particles propagating over large distances ($X \gg P/m^2$) are constrained by the conservation laws at the vertex to be on shell. The on-shell condition is all that is needed to fix the phase difference and the oscillation length.

Consider a diagram with a propagator between vertices at spacetime points (x_1, x_2) with a number of external legs at the vertices. The amplitude is proportional to

$$\int d^4x_1 d^4x_2 \exp(-i \sum_i q_i \cdot x_1) K(x_1, x_2) \exp(i \sum_f q_f \cdot x_2) \quad (3.1)$$

Where q_i are the incoming four momenta from the external legs at x_1 and q_f are the incoming four momenta from the external legs at x_2 . Substituting the propagator

$$K(x_1, x_2) = \int d^4p \frac{e^{-ip \cdot (x_1 - x_2)}}{p^2 - m^2 + i\epsilon} \quad (3.2)$$

in (3.1) and integrating over x_1 and x_2 gives the usual energy momentum conservation $\delta^4(\sum_i q_i - p)$ and $\delta^4(\sum_f q_f - p)$ at the vertices. When the proper distance between the two vertices s is larger than m^{-1} , then the propagator used in (3.1) is the asymptotic form (2.7), which in momentum space can be written as [17],

$$\begin{aligned} K(x_1, x_2; s \gg m^{-1}) &= \frac{1}{(2ms)^{3/2}} e^{-ims} \\ &= \int \frac{d^4p}{\sqrt{\mathbf{P}^2 + m^2}} \exp \{ -i(\sqrt{\mathbf{P}^2 + m^2})(t_f - t_i) + i\mathbf{P} \cdot (\mathbf{x}_f - \mathbf{x}_i) \} \end{aligned} \quad (3.3)$$

Substituting (3.3) in (3.1) and integrating over x_1 we obtain

$$\begin{aligned} \int d^4x_1 \exp \{ -i(\sum_i E_i - \sqrt{\mathbf{P}^2 + m^2})t_1 + i(\sum_i \mathbf{q}_i - \mathbf{P}) \cdot \mathbf{x}_1 \} \\ = \delta(\sum_i E_i - \sqrt{\mathbf{P}^2 + m^2}) \delta^3(\sum_i \mathbf{q}_i - \mathbf{P}) \end{aligned} \quad (3.4)$$

This implies that at a vertex energy and momentum are conserved and in addition those particles which propagate without freely over distances larger than $(|\mathbf{P}|/m^2)$ obey the mass shell constraint, $E = \sqrt{\mathbf{P}^2 + m^2}$.

4 Path Integral method

From the derivation of the propagation amplitude (2.7) it is clear that there is no difference in the result on whether the propagating fields are bosons or fermions. The same result for the propagation amplitude is therefore also expected in relativistic quantum mechanics. In relativistic quantum mechanics the propagation amplitude $K(m_i, t)$ is computed from the classical action for a relativistic particle by the path integral method [15]. The amplitude for a particle of mass m_i to propagate from the spacetime point x_i to another spacetime point x_f can be written formally as,

$$K(x_f, x_i; m) \equiv \int \mathcal{D}[x(\tau)] \exp \left\{ -\frac{im}{2} \int_{x_i}^{x_f} \left(\eta_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} + 1 \right) d\tau \right\} \quad (4.1)$$

For infinitesimal propagation between points x_j and x_{j+1} the quantum mechanical transition amplitude may be written as

$$K(x_{j+1}, x_j) = C \exp \left\{ -\frac{im}{2} \left(\frac{(x_{j+1} - x_j)^2}{(\Delta\tau)^2} + 1 \right) \Delta\tau \right\} \quad (4.2)$$

The condition $K(x_{j+1}, x_j) = \delta^4(x_{j+1} - x_j)$ as $\Delta\tau \rightarrow 0$ fixes the normalisation constant $C = (m/2\pi i \Delta\tau)^2$. In what follows we will not write out the explicit form for the normalisation factors as the final normalisation can be fixed from the boundary conditions.

To carry out the path integral we divide each path into $N \rightarrow \infty$ infinitesimal segments of length $\Delta\tau \rightarrow 0$ such that $N\Delta\tau = \tau$ the total path length. The amplitude for propagation from x_i to x_f is expressed as a product of the infinitesimal amplitudes $K(x_j, x_{j-1})$ over the small straight line segments $\Delta\tau$,

$$K(x_f, x_i) = \int dx_1 \cdots dx_{N-1} K(x_f, x_{N-1}) \cdots K(x_1, x_i) \quad (4.3)$$

By substituting the infinitesimal amplitudes (4.2) in the expression (4.3) we have the amplitude $K(x_f, x_i)$ as a product of ordinary integrals of the infinitesimal amplitudes $K(x_j, x_{j-1})$ over the intermediate spacetime points $(x_1, \cdots x_{N-1})$.

$$K(x_f, x_i) = C' \int dx_1 \cdots dx_{N-1} \exp \left\{ \frac{-im}{2\Delta\tau} \left(\sum_{j=1}^N (x_j - x_{j+1})^2 + N\Delta\tau \right) \right\} \quad (4.4)$$

The Gaussian integrals in (4.4) can be performed sequentially and the result of the $N-1$ integrations is

$$\int dx_1 \cdots dx_{N-1} \exp \left\{ \frac{-im}{2\Delta\tau} \left(\sum_{j=1}^N (x_j - x_{j+1})^2 \right) \right\} = C_N \exp \left\{ \left(\frac{-im}{2\Delta\tau} \right) \left(\frac{1}{N} (x_i - x_f)^2 \right) \right\} \quad (4.5)$$

Using (4.5) in (4.4) and writing $N\Delta\tau = \tau$ in the limit $N \rightarrow \infty$ and $\Delta\tau \rightarrow 0$, we have the expression for the propagation amplitude $K(x_i, x_f)$ as a function of the proper time of τ along some path,

$$K(x_i, x_f, \tau) = \left(\frac{m}{2\pi\tau} \right)^2 \exp \left\{ \frac{-im}{2} \left(\frac{(x_i - x_f)^2}{\tau} + \tau \right) \right\} \quad (4.6)$$

where the normalisation constant has been fixed by the requirement that $K(x_i, x_f, \tau) = \delta^4(x_i - x_f)$ as $\tau \rightarrow 0$.

In relativistic quantum mechanics the proper time along a path τ is an extra parameter that must be integrated over to include all paths including those which go backwards in the coordinate time t . The relativistic propagation amplitude from (4.6) is

$$K(x_f, x_i) = \int_0^\infty d\tau \left(\frac{m}{2\pi\tau} \right)^2 \exp \frac{-im}{2} \left(\frac{(x_i - x_f)^2}{\tau} + \tau \right) \quad (4.7)$$

The integral over τ can be performed by using the formula (2.4) to give the amplitude (4.7) as a Bessel function

$$K(x_f, x_i) = \left(\frac{m}{2\pi i} \right)^2 \frac{2}{s} K_1(ims) \quad (4.8)$$

which is of the same form as the field theory propagator (2.5) its asymptotic form

$$K(x_f, x_i; s \gg m^{-1}) \simeq \left(\frac{m}{2\pi s} \right)^{3/2} \exp(-ims) \quad (4.9)$$

is the same as (2.7).

The expression given in (4.8) is the correct form of the propagator for particles which in general are not on-shell. To compute the propagator for strictly on-shell (real) particles, we and restrict

ourselves only to those paths which go forward in time and propagate over a fixed total proper time $\tau = s = \sqrt{(x_f - x_i)^2}$. The on-shell propagator is then obtained by inserting $(2\pi\tau/m)^{1/2} \delta(\tau - \sqrt{(x_f - x_i)^2})$ (the factor of $(2\pi\tau/m)^{1/2}$ is for normalisation) in (4.7) to give,

$$K(x_f, x_i)_{on-shell} \simeq \left(\frac{m}{2\pi i s} \right)^{3/2} \exp(-ims) \quad (4.10)$$

Comparing (4.10) and (4.9) we see that in the asymptotic limit the off-shell propagator becomes identical to the on-shell propagator. In other words ‘virtual’ particles become ‘real’ on large distance propagation.

In non-relativistic quantum mechanics the proper time $\tau = (t_f - t_i)$ the coordinate time interval. The non-relativistic propagator (which off-course is always on-shell) is obtained by inserting $(2\pi\tau/m)^{1/2} \delta(\tau - (t_f - t_i)^2)$ in (4.7). The non-relativistic propagator thus obtained is the familiar form [15],

$$K_{NR}(\mathbf{x}_i, t_i; \mathbf{x}_f, t_f) = \left(\frac{m}{2\pi i(t_f - t_i)} \right)^{3/2} \exp\left\{-im(t_f - t_i) \left(1 - \frac{(\mathbf{x}_f - \mathbf{x}_i)^2}{2(t_f - t_i)^2}\right)\right\} \quad (4.11)$$

The amplitude (4.11) has the same form as the non-relativistic limit of the asymptotic propagator (2.7). As discussed earlier the phase factor $\exp - imt$ implies that in interference between two different mass states of non-relativistic particles (like K_L and K_S) the oscillations are periodic in $(\Delta m)t$.

5 Wavepacket analysis

The propagator $K(x_f, x_i)$ given in (2.7) is the amplitude for the propagation of a particle localised at x_i (a delta function initial wave function) to be detected at x_f . The propagator of some general initial wavepacket $\Psi_{in}(x - x_i)$ is obtained by using the expressions for the delta function propagator given in (2.7) and the superposition principle,

$$\tilde{K}(x_f - x_i) = \int d^4x K(x_f, x) \Psi_{in}(x - x_i) \quad (5.1)$$

If the initial wave function is a gaussian ,

$$\Psi_{in}(x - x_i) = N \exp\{-i(E_a(t - t_i) + i\mathbf{P}_a \cdot (\mathbf{x} - \mathbf{x}_i))\} \exp\left\{-\frac{(\mathbf{x} - \mathbf{x}_i)^2}{4\sigma_x^2} - \frac{(t - t_i)^2}{4\sigma_t^2}\right\} \quad (5.2)$$

where σ_x and σ_t are the uncertainties in the initial position and time of production of the particle respectively and N is the normalisation constant. In the earlier wave-packet analyses of oscillation problem [8], the uncertainty in the time of production was neglected. In a covariant treatment both should be included. In the next section we show that the time uncertainty gives rise to a novel phenomenon of conversion without oscillations in many experimentally relevant situations. Substituting the expression for $K(x_f, x)$ given in (2.7) and the gaussian initial wavepacket (5.2) in (5.1) and evaluating the integral by the stationary phase method [18] we obtain the expression for the propagation amplitude of a gaussian wavepacket ,

$$\tilde{K}(\mathbf{X}, T; m_a) = \left(\frac{1}{4\pi|\mathbf{X}|^2} \right)^{1/2} \left(\frac{\pi}{\sigma_x^2 + v_a^2 \sigma_t^2} \right)^{-1/4} \exp\left\{-i \frac{m_a^2}{E_a} T + i\mathbf{P}_a \cdot (\mathbf{X} - \mathbf{v}_a T)\right\} \exp\left\{-\frac{(\mathbf{X} - \mathbf{v}_a T)^2}{4(\sigma_x^2 + v_a^2 \sigma_t^2)}\right\} \quad (5.3)$$

where $\mathbf{X} = \mathbf{x}_f - \mathbf{x}_i$ and $T = t_f - t_i$ are the space and time intervals propagated by the center of the gaussian wavepacket and $\mathbf{v}_a \equiv \mathbf{P}_a/E_a$. The normalisation constant has been fixed such that $\int dT d^3\mathbf{X} \tilde{K}(\mathbf{X}, T; m_a) \tilde{K}^\dagger(\mathbf{X}, T; m_b) = 1$. The factor of $(1/4\pi|\mathbf{X}|^2)^{1/2}$ in (5.3) shows that the particle flux decreases inversely with the square of distance. Neutrino disappearance experiments look for evidence of depletion of a certain neutrino species over and above the expected inverse square decrease in flux. In the following we will not display this factor $(1/4\pi|\mathbf{X}|^2)$ in the probability expressions. The amplitude for the oscillation of one type of neutrino flavour to another is obtained by substituting the propagation amplitudes for mass eigenstates (5.3) in ,

$$\mathcal{A}(\alpha \rightarrow \beta; X, T) = \sum_a^3 U_{\alpha a} \tilde{K}(m_a; X, T) U_{a\beta}^* \quad (5.4)$$

where α, β denote the flavor eigenstates (ν_e, ν_μ, ν_τ or $K^0, \bar{K}^0$) and the summation index a denotes a mass eigenstates (ν_1, ν_2, ν_3 or K_L, K_S). $U_{\alpha a} = \langle \alpha | a \rangle$ and $U_{\beta a}^* = \langle a | \beta \rangle$ are the elements of the mixing matrix which relate the flavor eigenstates with the mass eigenstates. The probability of flavour oscillation as a function of space-time is the modulus squared of the amplitude (1.4)

$$\begin{aligned} P(\alpha \rightarrow \beta; X, T) &= \left| \sum_a^3 U_{\alpha a} \tilde{K}(m_a; X, T) U_{a\beta}^* \right|^2 \\ &= \sum_a |U_{\beta a}|^2 |U_{\alpha a}|^2 + \sum_{a \neq b} U_{\beta a} U_{\alpha a}^* U_{\beta b}^* U_{\alpha b} \tilde{K}(\mathbf{X}, T; m_a) \tilde{K}^\dagger(\mathbf{X}, T; m_b) \end{aligned} \quad (5.5)$$

In interference experiments the time of flight of the particle is not measured and only the distance between the source and the detector is accurately known [16]. The probability for the flavour conversion as a function of distance is given by the time integral of (5.5) ,

$$\begin{aligned} P(\alpha \rightarrow \beta; X) &= \int dT \left| \sum_a^3 U_{\alpha a} \tilde{K}(m_a; X, T) U_{a\beta}^* \right|^2 \\ &= \sum_a |U_{\beta a}|^2 |U_{\alpha a}|^2 + \sum_{a \neq b} U_{\beta a} U_{\alpha a}^* U_{\beta b}^* U_{\alpha b} \int dT \tilde{K}(\mathbf{X}, T; m_a) \tilde{K}^\dagger(\mathbf{X}, T; m_b) \end{aligned} \quad (5.6)$$

The interference term given by the time-overlap of the propagation amplitudes (wave-functions) of different mass eigenstates can be evaluated for the gaussian propagator (5.3) and is given by,

$$\begin{aligned} \text{Re} \int dT \tilde{K}(\mathbf{X}, T; m_a) \tilde{K}^\dagger(\mathbf{X}, T; m_b) &= \frac{2}{v} \cos\{(\mathbf{P}_a - \mathbf{P}_b) \cdot \mathbf{X} - (E_a - E_b) \frac{(\mathbf{v}_a + \mathbf{v}_b) \cdot \mathbf{X}}{2v^2}\} \\ &\times \exp\left\{- (E_a - E_b)^2 \frac{(\sigma_x^2 + v^2 \sigma_t^2)}{2v^2} - \frac{(\mathbf{v}_a - \mathbf{v}_b)^2 X^2}{8(\sigma_x^2 + v^2 \sigma_t^2)v^2}\right\} \end{aligned} \quad (5.7)$$

where $v \equiv \sqrt{(v_a^2 + v_b^2)/2}$. In terms of the average momentum $\mathbf{P}$ and energy E and their respective differences $\Delta\mathbf{P}$ and ΔE the interference term (5.7) ,to the leading order in $(\Delta P/P)$ and $(\Delta E/E)$, reduces to the form,

$$\frac{2}{v} \cos\left\{\frac{X}{P}(E\Delta E - \mathbf{P} \cdot \Delta\mathbf{P})\right\} \exp\{-A\} \quad (5.8)$$

where the exponential damping factor,

$$A = (\Delta E)^2 \frac{2\bar{\sigma}^2}{P^2} + \left(\frac{\Delta P}{P}\right)^2 \frac{X^2}{4\bar{\sigma}^2} \quad (5.9)$$

and $\bar{\sigma}^2 \equiv (\sigma_x^2 + (P/E)^2 \sigma_t^2)$.

In general neither ΔE nor ΔP is zero and they depend upon how the state is prepared. For example if the momentum of the initial and the associated final state is measured then $\Delta P = 0$. In the last section we have shown that for long distance propagation ($s > m^{-1}$) $\mathbf{P}$ and E are not independent and are related by the mass shell constraint. The particular combination that appears in the phase difference turns out to be independent of the preparation and is fixed by the condition that each of the mass eigenstates be on shell,

$$\begin{aligned} E_i^2 - P_i^2 &= m_i^2 \quad i = a, b \\ \Rightarrow 2E\Delta E - 2\mathbf{P} \cdot \Delta\mathbf{P} &= \Delta m^2 \end{aligned} \quad (5.10)$$

Substituting (5.10) in (5.8) we see that the interference term of (5.7) is given by ,

$$\frac{2}{v} \cos\left(\frac{\Delta m^2}{2P} X\right) e^{-A} \quad , \quad (5.11)$$

The probability for the flavour conversion as a function of distance (5.6) is therefore,

$$\begin{aligned} P(\alpha \rightarrow \beta; X) &= \sum_a \frac{1}{v_a} |U_{\beta a}|^2 |U_{\alpha a}|^2 \\ &+ \sum_{a \neq b} \frac{1}{v} |U_{\beta a} U_{\alpha a}^* U_{\beta b}^* U_{\alpha b}| \cos\left(\frac{(m_a^2 - m_b^2)}{2P} X + \delta\right) e^{-A} \end{aligned} \quad (5.12)$$

where $\delta = \arg(U_{\beta a} U_{\alpha a}^* U_{\beta b}^* U_{\alpha b})$.

We see that the standard oscillation formula which was obtained for relativistic particles [1] is actually valid at all energies. In other words although the formula (5.12) is usually derived by taking the leading term in a (m^2/P^2) series, our covariant calculation shows that there are actually no $O(m^2/P^2)$ corrections to (5.12).

In the non-relativistic regime where $P = (m_a + m_b)v/2$, the standard kaon oscillation formula $\cos(m_a - m_b)T$ is recovered.

There is no extra factor of two in the relativistic kaon oscillation formula as would have appeared without the wavepacket averaging . We see that although the two mass eigenstates have different proper times the wave packet overlap results in the average proper time appearing in the interference term. Therefore the interference phase is actually $(m_a - m_b)\bar{s} = (m_a - m_b)(m_a + m_b)T/(E_a + E_b) = (m_a^2 - m_b^2)T/2E$. The average time prescription of [5] can therefore be justified using the covariant propagator method. Other methods of showing that an extra factor of two does not appear in the flavour oscillations formula have been discussed in [6].

6 No muon oscillation

The energy momentum relation was employed in the last section to express the oscillation length in terms of the average momentum of the mass eigenstates. We can use the conservation laws to give an explicit expression for the phase difference of the oscillation term. For example if the neutrinos are produced by pion decay $\pi^+ \rightarrow \mu^+ \nu_\mu$, one can use the conservation laws to write the oscillation length of the neutrino in terms of m_π and m_μ and explicitly the validity of the formula for the phase difference (5.11). In the pion rest frame the energy and the momentum of the neutrino and the muon are,

$$\begin{aligned} E_\nu &= \frac{m_\pi^2 - m_\mu^2 + m_\nu^2}{2m_\pi} \\ E_\mu &= \frac{m_\pi^2 + m_\mu^2 - m_\nu^2}{2m_\pi} \\ P &= |\mathbf{P}_\nu| = |\mathbf{P}_\mu| = \frac{((m_\pi^2 - (m_\nu + m_\mu)^2)(m_\pi^2 - (m_\nu - m_\mu)^2))^{1/2}}{2m_\pi} \end{aligned} \quad (6.1)$$

Using (6.1) we can check explicitly the expression for the neutrino phase difference,

$$\begin{aligned} \Delta E_\nu &= \frac{\Delta m_\nu^2}{2m_\pi} \\ \Delta P_\nu &= \frac{\Delta m_\nu^2}{2m_\pi P} (E_\nu - m_\pi) \\ \Delta \phi_\nu &= (E_\nu \Delta E_\nu - \mathbf{P}_\nu \cdot \Delta \mathbf{P}_\nu) \frac{X}{P_\nu} = \frac{\Delta m_\nu^2}{2P} X \end{aligned} \quad (6.2)$$

Another application of the conservation laws is the question of associated particle oscillation raised by SWS [4]. For instance in the process $\pi^+ \rightarrow \mu^+ \nu_\mu$, SWS state that since the muon momenta depend on m_ν , an interference of muon eigenstates corresponding to different $P_\mu(m_\nu)$ will result in the occurrence of oscillation in space of the muon probability. Similarly in the reaction $p p \rightarrow \Lambda K$, SWS claim that the associated oscillations of Λ probability in space ought to be seen. This claim has been disproved by Lowe et al and Okun et al, by different methods. We will see that the phase difference of m_μ corresponding to the different ν mass states actually vanishes. Using (6.1) we see that

$$\begin{aligned} \Delta E_\mu &= -\frac{\Delta m_\nu^2}{2m_\pi} \\ \Delta P_\mu &= -\frac{\Delta m_\nu^2}{2m_\pi P} E_\mu \\ \Delta \phi_\mu &= (E_\mu \Delta E_\mu - \mathbf{P}_\mu \cdot \Delta \mathbf{P}_\mu) \frac{X}{P_\mu} = 0. \end{aligned} \quad (6.3)$$

Okun et al state that muon oscillations do not take place, if the neutrino is not observed 'owing to the orthogonality of the neutrino eigenstates. That would imply that if there is a mixing of the neutrinos with some heavy states, which would make the low-lying mass neutrinos non-orthogonal,

then muon oscillations will take place. This would have been powerful method for testing for the orthogonality of the observed neutrinos. However from (6.3) we see that the form of the phase difference is such that the conservation laws ensure that whether the ν is measured or not there is no EPR correlation to the μ phase difference. There is an EPR type correlation of E_μ and P_μ with the neutrino mass eigenstates but the phase difference vanishes. Therefore there will be no muon oscillations In the linear combination of different mass eigenstates, neither ΔE nor ΔP is zero. Assuming that either of them is zero would lead to incorrect results.

7 Conversion without oscillation

The suppression factor A which goes with the oscillation term has some interesting new implications. If the parameters of the experiment are such that A becomes large then no oscillations is spacetime can be observed. What can be observed is a constant conversion probability.

For example, the survival probability of a ν_μ after propagating over a distance X in the large A limit is, (for two neutrino flavours (ν_e, ν_μ) with the mixing matrix elements $U_{e1} = U_{\mu 2} = \cos\theta$ and $U_{e2} = -U_{\mu 1} = \sin\theta$),

$$P(\nu_\mu \rightarrow \nu_\mu; X) = \sin^4\theta + \cos^4\theta + 2\sin^2\theta\cos^2\theta \cos(\frac{\Delta m^2}{2P}X) e^{-A} \simeq \sin^4\theta + \cos^4\theta \quad (7.1)$$

and the expression for the conversion probability when A is large is,

$$P(\nu_\mu \rightarrow \nu_e; t) = 2\sin^2\theta \cos^2\theta (1 - \cos(\frac{\Delta m^2}{2P}X) e^{-A}) \simeq 2\sin^2\theta \cos^2\theta \quad (7.2)$$

It has been noted earlier [8] one condition A must be small which implies that X must be smaller than the coherence length $L_{coh} = (4\sqrt{2}\bar{\sigma}P^2/\Delta m^2)$. We shall show below that the constraint $L_{osc} < L_{coh}$ is not a sufficient condition to ensure the occurrence of flavour oscillations in space. In the analysis of refs [8] only the uncertainty in the initial position σ_x was considered, in our analysis we have included the contribution of σ_t the uncertainty in time at which the neutrino was produced, which as it turns out, makes the larger contribution to the suppression term A . This happens when the neutrinos are produced from long lived resonances. The neutrino wavepacket which is produced has a spread in time with width σ_t which cannot be smaller than the lifetime of the resonance whose decay produces it. The dominant contribution in that case arises from the first term of A ,

$$A \simeq (\Delta E)^2 \frac{\bar{\sigma}^2}{2} \simeq (\frac{\Delta m^2 \tau}{2\sqrt{2}E})^2 \quad (7.3)$$

where we have equated the initial time uncertainty σ_t with τ - the lifetime of the resonance that produces the neutrino, and we have ignored the spatial spread of the wave-packet σ_x which in most experiments is many orders of magnitude smaller than σ_t . In terms of the oscillation length $L_{osc} = 4\pi E/(\Delta m^2)$ the suppression factor A . In order to observe oscillations the source-detector distance L must be larger than L_{osc} . Which that the minimum source-detector distance in order that neutrino oscillations be observed is given by condition, $L_{min} = (\pi)\sqrt{2}\tau$. We shall show below that most neutrino experiments do not satisfy this criterion for observability of space oscillations. On the other hand spatial oscillations of flavour can be observed in the Φ and B factories owing to the large width of ϕ and Υ resonances.

For relativistic particles produced from long lived resonances the formula for the conversion probability which should be fitted with the experiments is ,

$$P(\nu_\alpha \rightarrow \nu_\beta) = \frac{1}{2} \sin^2 2\theta \left(1 - \cos\left(\frac{2.53\Delta m^2 L}{E}\right) \exp\left\{-\left(\frac{1.79\Delta m^2 \tau}{E}\right)^2\right\} \right) \quad (7.4)$$

where Δm^2 is the mass square difference in eV^2 , L is the detector distance in m (km), τ is the lifetime of the parent particle in the lab frame in m (km) and E is the energy in MeV (GeV).

8 Constraints on Δm^2 and $\sin^2 2\theta$ from experiments

We shall now fit the oscillation formula (7.4) with the results of some terrestrial neutrino oscillation experiments , classified according to the type of the source of neutrinos.

Stopped Muons : The LSND experiment [9] has searched for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ in a beam of anti-muon neutrinos produced from stopped muons. The minimum uncertainty in the time of $\bar{\nu}_\mu$ production is the muon lifetime $\tau_\mu = 658.65m$. The average neutrino energy $E_\nu \simeq 30MeV$ and the detector is at a distance $L = 30m$ from the source. In order that neutrino oscillations from muon decay neutrinos be seen the minimum source detector distance must be $L_{min} = \sqrt{2}\pi\tau_\mu = 2927.6m$. This means that the conversion probability does not oscillate as a function of distance. Fitting the reported [9] range of conversion probability $P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) = (0.31 + 0.11 - 0.1 \pm 0.05) \times 10^{-2}$ with the formula (7.4) , the regions of allowed Δm^2 and $\sin^2 2\theta$ which gives rise to this probability band are shown as dotted band in Fig. 1. The same range of conversion probability when fitted with the standard oscillation formula gives the shaded region shown in Fig.1. We have averaged the conversion probability by assuming a gaussian distribution of neutrino energy, with average energy $30MeV$ and a half width of $10MeV$. The asymptotic bounds are as follows. If $\Delta m^2 > (2\sqrt{2}E/\tau_\mu) = 2.54 \times 10^{-2}eV^2$ then mixing angle constrained in the range $0.3 \times 10^{-2} \leq \sin^2 2\theta \leq 0.9 \times 10^{-2}$. For large mixing angles $\sin^2 2\theta \simeq 1$ the mass square difference lies in the range, $2 \times 10^{-3}eV^2 \leq \Delta m^2 \leq 3 \times 10^{-3}eV^2$. This bound is two orders lower than what is obtained by the use the standard oscillation formula [9].

Stopped Pions : The Karmen experiment [11] searches for $\nu_\mu \rightarrow \nu_e$ conversion in a beam of ν_μ produced by the decay of stopped pions . The time at which the ν_μ is prduced cannot be localised to less than the pion lifetime $\tau_{\pi^+} = 7.804m$. The ν_μ energy is $E_\nu = 29.8MeV$. In order for oscillations to be observable the detector distance must be larger than $L = \sqrt{2}\pi\tau_{\pi^+} = 35m$. Since the detector distance is $17.5m$, it is not possible to observe spatial oscillations in the Karmen experiment. The experimental bound $P(\nu_\mu \rightarrow \nu_e) < 3 \times 10^{-3}$, fitted with the conversion probabiltiy formula (7.4) constrains $\sin^2 2\theta < 0.6 \times 10^{-2}$ if $\Delta m^2 > (2\sqrt{2}E/\tau_{\pi^+}) = 2 eV^2$. In the $\sin^2 2\theta \sim 1$ limit $\Delta m^2 < 0.16eV^2$. We have averaged the conversion probability by assuming a gaussian distribution of neutrino energy, with average energy $30MeV$ and a half width of $10MeV$. The regions of the parameter space allowed by the covariant wavepacket formula and by the standard formula are shown in Fig.2. Since the pion decay length is of the same order as the experimental baseline, the improvement in sensitivity to Δm^2 is marginal compared to LSND.

High energy Pions or Kaons: When high energy neutrino beams are obtained from the decay of a pion or kaon in flight the both τ and E are larger by the Lorentz factor γ and consequently A remains the same as that with stopped pions or kaons. In the BNL E776 experiment [10] most of the neutrinos are produced in $\pi^\pm$ decay and have an average energy $E \simeq 5GeV$ and the detector is at a distance $L = 1km$. In the lab frame the neutrino time uncertainty is $\tau = \tau_{\pi^+}(E_{\pi^+}/m_{\pi^+})$. Taking the

mean $\pi^\pm$ energy to be 10GeV , the lifetime of the pions in the lab frame is $\tau = 7.804(10/0.135)m = 0.578\text{km}$. Fitting the E776 experimental limit $P(\nu_\mu \rightarrow \nu_e) < 1.5 \times 10^{-3}$ with the conversion probability formula (7.4), we see from Fig.3 that $\sin^2 2\theta < 0.3 \times 10^{-2}$ if $\Delta m^2 > (2\sqrt{2}E/\tau_{\pi^\pm}) = 5 \text{ eV}^2$. In the $\sin^2 2\theta \sim 1$ limit, $\Delta m^2 < 0.2\text{eV}^2$.

In the CCFR [12] experiments most of the neutrinos are produced in $K^\pm$ decay and have an average energy $E \simeq 140\text{GeV}$ and the detector is at a distance $L = 1.4\text{km}$. In the lab frame the neutrino time uncertainty is $\tau = \tau_{K^\pm}(E_{K^\pm}/m_{K^\pm})$. Taking the mean $K^\pm$ energy to be 600GeV we see that the lifetime of the kaons in the lab frame is $\tau = 3.7(600/0.493)m = 4.514\text{km}$. Fitting the CCFR experimental limit $P(\nu_\mu \rightarrow \nu_e) < 0.9 \times 10^{-3}$ with the conversion probability formula (7.4) we see from Fig.3 that $\sin^2 2\theta < 1.8 \times 10^{-3}$ if $\Delta m^2 > (2\sqrt{2}E/\tau_{K^\pm}) = 76 \text{ eV}^2$. In the $\sin^2 2\theta \sim 1$ limit, $\Delta m^2 < 1.2\text{eV}^2$.

Reactor neutrinos: Reactor neutrinos are produced primarily from the beta decay of ^{235}U and ^{239}Pu which have a half life $\sim 10^2\text{sec}$ [13]. The minimum detector distance for observing neutrino oscillations is 10^{11}m which means that it is impossible to detect spatial oscillations with reactor neutrinos. The upper bounds on conversion probability in the Bugey experiment is $P(\bar{\nu}_e \rightarrow \bar{\nu}_\mu \text{ or } \bar{\nu}_\tau) < 0.075$. The largest detector distance $L = 95\text{m}$ and the average neutrino energy is $E \simeq 5\text{MeV}$. The conversion probability formula is independent of Δm^2 if $\Delta m^2 > 10^{-9}\text{eV}^2$. Fig 3. shows that in this limit the reactor experiments rule out $\sin^2 2\theta \geq 0.15$. In the standard oscillation formula the oscillatory term averages to zero when $\Delta m^2 > (4\pi E/L)$. Using the wavepacket formula (7.4) however we see that the oscillatory term is exponentially damped at much lower values of Δm^2 . If neutrino sources have a large decay time which is the case in most experiments, the conversion probability is sensitive to mass differences $\Delta m^2 > (2\sqrt{2}E/\tau)$. For this reason the LSND experiment which uses neutrinos from muon decay has two orders of magnitude more sensitive than KARMEN, BNL-E776 etc where the neutrinos are from π and K decay. This is evident from the combined plot of these experimental results, shown in Fig.3 fitted with the covariant wavepacket formula (7.4). The reactor experiments where $\tau \sim 10^2\text{s}$ are in fact sensitive to values of Δm^2 as low as 10^{-9}eV^2 . But the probability measurement in reactor experiments is poor compared to the accelerator experiments which is why they rule out only a small region of parameter space.

Meson oscillations at B and Φ factories: The mesons produced by ϕ or $\Upsilon(4S)$ decay are non-relativistic and the expression for the damping factor is $A = 2(\Delta m \tau)^2$. For the kaons produced from ϕ decays, $\Delta m_K = 3.5 \times 10^{-12}\text{MeV}$ and the ϕ lifetime is $\tau_\phi = 0.225\text{MeV}^{-1}$, and the damping factor $A = 2.4 \times 10^{-24}$ is negligible. For the case of $B\bar{B}$ oscillations produced from $\Upsilon(4S)$ decays, the $\Delta m_B = 3.53 \times 10^{-10} \text{ MeV}$ and the lifetime of $\Upsilon(4S)$ is $\tau_\Upsilon = 0.042\text{MeV}^{-1}$ and again the damping factor, $A = 8.6 \times 10^{-22}$ is negligible. This is important for tests of EPR [19] and Quantum Mechanics [20] where the measurement of the oscillatory term of the probability is essential.

9 Conclusions

The covariant formulation gives two new results compared to the standard treatment. We use the condition that the spread of the neutrino wave function along the time axis is given by the lifetime of the particle whose decay produces the neutrino. When neutrinos are produced from long lived particles like muons, the interference term in the conversion probability formula vanishes for much smaller values of the neutrino mass square difference compared with the standard formula. The

reason this happens can be understood with the following picture. When the spread of the neutrino is large along the time axis, its spread along the energy axis is small. The neutrino wavefunctions for different masses are then energy eigenstates which are mutually orthogonal. The interference term which is the overlap of the wavefunctions of different mass states therefore vanishes when these states become orthogonal.

The second new result is that for the secondary particles, like muons in the decay $\pi \rightarrow \mu\nu$ there is no spacetime variation of probability independent of whether the neutrinos are measured or not. Although the muon wavefunction is a linear combination of distinct energy and momentum states corresponding to the different ν mass states, the phase difference for the secondary particles vanishes. Therefore there is no EPR type oscillation of the secondary particles probability distribution. This is true independent of whether the primary oscillations of the ν 's is measured or not. This conclusion results from the application of the correct conservation laws according to which the different mass states of ν have neither energy nor momentum in common as is sometime assumed in the standard derivations.

Acknowledgments I thank Terry Goldman, Walter Grimus and Marek Nowakowski for useful correspondence.




Figure 1: Lsnu μ decay at rest experiment [9] allowed regions with the standard formula (between the dotted curves) and the covariant oscillation formula (continuous curves).




Figure 2: Karmen π decay at rest experiment [11] allowed regions with the standard formula (dotted line) and the covariant oscillation formulas (dashed line).




Figure 3: The region between the continuous lines is allowed by the LSND μ decay in flight experiment [9]. The region between the dotted lines is allowed by the LSND π decay in flight experiment [21]. Region ruled out by E776 [10] is above the dashed-dotted curve and by Karmen [11] is above dashed curve. Region above the topmost dashed curve is ruled out by CCFR [12].

References

- [1] B. Pontecorvo, Sov. Phys. JETP **26**, 984 (1968);
S. M. Bilenky and S. T. Petcov, Rev. Mod. Phys. **59**, 671 (1987);
R. N. Mohapatra and P. B. Pal, *Massive Neutrinos in Physics and Astrophysics* (World Scientific, Singapore, 1991);
C. W. Kim and A. Pevsner, *Neutrinos in Physics and Astrophysics*, harwood academic publishers, Chur,(1993).
- [2] E.D.Cummins and P.H.Bucksbaum, *Weak interactions of leptons and quarks*, Cambridge University Press,(1983);
R.P.Feynman , R.B.Leighton and M.Sands *Feynman Lectures in Physics Vol III*, Addison Wesley, Reading MA (1965).
- [3] Kayser and Stodolsky, Phys. Lett.**B359**, 333(1995).
- [4] Y.N.Srivastava, A.Widom and E.Sassaroli, Z. fur Physik, **C66**,601 (1995).
- [5] J.Lowe, B. Bassallek, H. Burkhardt, A.Rusek, G.J.Stephenson Jr and T.Goldman , Phys. Lett **B384**, 228(1996) ;
B.Kaysor, hep-ph 9702327.
- [6] Yu. Grossman and H.J.Lipkin, Phys Rev **D55**, 2760 (1996);
B.Anchoea, A.Bramon, R.Muniz-Tapia and M. Nowakowski , Phys Lett **B389**, 149(1996) ;
A.Dolgov, A.Yu.Morozov, L.B.Okun and M.G.Schepkin hep-ph 9703241;
M.M. Nieto, hep-ph 9509370 ;
T. Goldman , hep-ph 9604357.
- [7] W.Grimus and P. Stockinger, Phys Rev **D54**, 3414 ,(1996).
- [8] C. Giunti, C. W. Kim, J. A. Lee and U. W. Lee, Phys. Rev. D **48**, 4310 (1993);
B. Kayser, Phys. Rev. D **24**, 110 (1981);
S.Nussinov, Phys Lett**B63** , 201 (1976).
- [9] C.Athanassopoulos et al (LSND) , Phys Rev Lett. **75** , 2650 (1995).
- [10] L.Brodovsky et al (BNL-E776), Phys. Rev Lett **68**, 274 (1992) .
- [11] B.Armbrusuter et al (Karmen), Nucl.Phys. **B38** ,235 (1995).
- [12] A. Romosan et al (CCFR), Phys. Rev. Letters ,**78**, 2912(1997).
- [13] B.Achkar et al (Bugey), Nucl. Phys. **B434** , 503 (1995).
- [14] I.S.GradshTEyn and I.M.Ryzhik, Academic Press, New York, p 956 , 8.427, (1981).
- [15] R.P.Feynman and A.R. Hibbs, *Quantum Mechanics and Path Integrals*, McGraw Hill, New York, (1965.).

- [16] H.J. Lipkin , Phys. Lett.**B348**, 604 (1995).
- [17] N.N. Bogoliubov and D.V. Shirkov, *An introduction to the theory of quantised fields* , Wiley Interscience, New York, (1959).
- [18] L. Mandel and E.Wolf, *Optical coherence and quantum optics*, p 128, Cambridge University Press, (1995).
- [19] H.J.Lipkin , Phys Rev **176**, 1715 (1968);
B.Kayser and L.Stodolsky, Phys Lett **359B**, 343 (1996).
- [20] J.Ellis,J.S.Hagelin,D.V. Nanoupoulos and M. Srednicki, Nucl Phys **241B**, 381(1984);
P.Huet and M.Peshkin, Nucl. Phys. **B 434**, 3 (1995);
R.A Bartleman and W.Grimus, Phys Lett **392B**, 426 (1997).
- [21] C.Athanassopoulos et al (LSND) nucl-ex/9706006